

# Improvements in Liver Fibroinflammation (as assessed by Corrected T1 [cT1]) with HTD1801 (Berberine Ursodeoxycholate) Treatment in Patients with Metabolic Dysfunction-Associated Steatohepatitis and Type 2 Diabetes Mellitus

Stephen A. Harrison<sup>1</sup>, Nadege Gunn<sup>2</sup>, Guy Neff<sup>3</sup>, Abigail Flyer<sup>4</sup>, Alexander Liberman<sup>5</sup>, Leigh MacConell<sup>5</sup>



## BACKGROUND

- HTD1801 (berberine ursodeoxycholate), a new molecular entity, is a gut-liver anti-inflammatory metabolic modulator
  - Targets multiple pathways pivotal to metabolic regulation, including those associated with metabolic and digestive diseases
- Corrected T1 (cT1) is an MRI-based quantitative metric for assessing liver inflammation and fibrosis
- Previous studies in patients with metabolic dysfunction-associated steatohepatitis (MASH, formerly nonalcoholic steatohepatitis) have reported that<sup>1</sup>:
  - cT1 improvements are moderately correlated with histologic improvements in NAFLD Activity Score (NAS) and fibrosis
  - cT1 levels are associated with clinical outcomes (liver and cardiovascular disease)
- In a Phase 2 study in patients with MASH and type 2 diabetes mellitus (T2DM), HTD1801 significantly reduced liver fat content (LFC) as determined by MRI-PDFF after 18 weeks of treatment (NCT03656744)<sup>2</sup>

The objective of this post-hoc analysis was to evaluate the effects of HTD1801 on cT1 in patients with MASH and T2DM

## METHODS

### Phase 2a Proof of Concept Dose-Finding Study<sup>2</sup>


#### Key Entry Criteria

- Presumed MASH with LFC ≥10%
- cT1 ≥830 ms
- Serum AST ≥20 U/L
- T2DM and on stable therapy

#### Primary Endpoint

Change from baseline in LFC by MRI-PDFF

- MRI-PDFF data was collected prospectively for evaluation of the primary endpoint
- cT1 segmented analysis was evaluated after study completion for HTD1801 1000 mg BID or placebo



## RESULTS



### Demographics and Baseline Characteristics

	Placebo (N=33)	HTD1801 1000 mg BID (N=34)
Age, years	58 (11)	53 (12)
Female, n (%)	22 (67)	24 (71)
White, n (%)	31 (94)	31 (91)
Hispanic or Latino, n (%)	13 (39)	11 (32)
BMI, kg/m <sup>2</sup>	35 (6)	36 (6)
HbA1c, %	7.0 (1)	7.3 (1)
MRI-PDFF, %	20 (6)	19 (7)
cT1, ms*	938 (98)	942 (91)
ALT, U/L	54 (27)	62 (32)

Values are Mean (SD) unless otherwise noted.

\*Baseline cT1 values were reassessed using a segmented analysis as defined by the analysis plan rather than the regional analysis used to determine subject eligibility at screening.


### Patients Receiving HTD1801 Had A Significant Reduction in Fibroinflammatory Disease as Assessed by cT1




\*p<0.05, p-values are obtained from an ANCOVA model with treatment group as a fixed effect, and baseline ALT and baseline cT1 as covariates.

## RESULTS

### More Patients Treated with HTD1801 Achieved Reductions in Biomarkers Correlated with Histologic Improvement<sup>1,4-5</sup>



### A Larger Proportion of Patients Receiving HTD1801 Achieved Clinically Meaningful Thresholds in cT1



\*p<0.05, 2-sided p-values are obtained from a chi-square test.

- cT1 <800 ms has been associated with a low likelihood of disease activity<sup>3</sup>
- A cT1 reduction of ≥80 ms has been correlated with improved histology (a 2-point reduction in NAS)<sup>1</sup>
- After 18 weeks:
  - Treatment with HTD1801 resulted in a larger proportion of patients achieving a cT1 <800 ms
  - Twice as many patients treated with HTD1801 compared to placebo achieved either a cT1 <800 ms or ≥80 ms reduction in cT1

### TEAEs Occurring in More Than 2 Subjects<sup>2</sup>

	Placebo (N=33)	HTD1801 1000 mg BID (N=34)
Any TEAE, n (%)	20 (61)	26 (76)
Diarrhea	3 (9)	11 (32)
Nausea	3 (9)	7 (21)
Headache	2 (6)	3 (9)
Upper Respiratory Tract Infection	4 (12)	1 (3)
Abdominal Pain	3 (9)	1 (3)

- The incidence of TEAEs was low and events were generally mild in severity
  - The most common TEAEs were GI-related events, which occurred more frequently with HTD1801
- Three serious adverse events occurred during the study none of which were considered related to study drug
  - Includes myocardial infarction (1000 mg BID); oxygen saturation decreased (500 mg BID); and bladder transitional cell carcinoma (placebo)

## SUMMARY

- Subjects receiving HTD1801 had significant reduction in fibroinflammatory disease as assessed by cT1
- Across multiple biomarkers, HTD1801 resulted in more patients achieving clinically relevant thresholds correlated with histologic improvement and lower disease activity
- These data suggest that HTD1801 may improve liver histology in patients with MASH and T2DM, warranting further investigation
- A Phase 2b study is currently ongoing to evaluate the histologic effects of HTD1801 in patients with MASH and T2DM or prediabetes (NCT05623189)

#### References

- Dennis A, et al. *Front Endocrinol.* 2021;11:575843.
- Harrison SA, et al. *Nat Commun.* 2021;12(1):5503.
- Andersson A, et al. *Clin Gastroenterol Hepatol.* 2022;20:2451-61.
- Stine J, et al. *Clin Gastroenterol Hepatol.* 2021;19(11):2274-283.
- Loomba R, et al. *Gastroenterology.* 2019;156(1):88-95.

#### Author Affiliations

- Pinnacle Clinical Research, TX, USA; 2. Impact Research Institute, TX, USA; 3. Covenant Metabolic Specialists, LLC, FL, USA;
- Pacific Northwest Statistical Consulting, WA, USA; 5. HighTide Therapeutics, MD, USA

Contact Information HighTide Therapeutics: info@hightidex.com